

Periodical Solutions of Multi-Time Hamilton Equations

Constantin Udrişte, Iulian Duca

Abstract

To our knowledge, there are two main references [9], [12] regarding the periodical solutions of multi-time Euler-Lagrange systems, even if the multi-time equations appeared in 1935, being introduced by de Donder. That is why, the central objective of this paper is to solve an open problem raised in [12]: what we can say about periodical solutions of multi-time Hamilton systems when the Hamiltonian is convex?

Section 1 recall well-known facts regarding the equivalence between Euler-Lagrange equations and Hamilton equations. Section 2 analyzes the action that produces multi-time Hamilton equations, and introduces the Legendre transform of a Hamiltonian together a new dual action. Section 3 proves the existence of periodical solutions of multi-time Hamilton equations via periodical extremals of the dual action, when the Hamiltonian is convex.

Key words: multi-time Hamilton action, periodical extremals, convex Hamiltonian.

2000 Mathematics Subject Classification: 53C21, 49K20, 49S05.

1 Classical Legendre transform of a Lagrangian

In this Section we recall the classical duality used in mechanics, based on the fact that the Legendre transform of a Lagrangian is a Hamiltonian (see [2], [7], [8], [11] for single-time theory, and [1], [3]-[6], [8]-[12] for multi-time theory).

1.1 Single-time Euler-Lagrange equations

The Lagrangian, $L : R \times R^{2n} \rightarrow R$, $(t, x, \dot{x}) \rightarrow L(t, x, \dot{x})$, where $x = (x^1(t), \dots, x^n(t))$ and $\dot{x} = (\dot{x}^1(t), \dots, \dot{x}^n(t))$, produces the single-time Euler-Lagrange equations

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}^i} = \frac{\partial L}{\partial x^i}, \quad i = 1, \dots, n. \quad (1)$$

1.2 Single-time Hamilton equations

We suppose that the Lagrangian L defines the diffeomorphism $\dot{x}^i \rightarrow p_i = \frac{\partial L}{\partial \dot{x}^i}$ called the *Legendre transformation*.

The Legendre transformation of the Lagrangian L is the Hamiltonian $H : R \times R^{2n} \rightarrow R$, $(t, x, p) \rightarrow H(t, x, p)$, $H(t, x, p) = \dot{x}^i p_i - L(t, x, \dot{x})$.

The equations (1) become

$$\frac{dx^i}{dt} = \frac{\partial H}{\partial p_i}, \quad \frac{dp_i}{dt} = -\frac{\partial H}{\partial x^i}, \quad i = 1, \dots, n \quad (2)$$

because

$$\frac{\partial H}{\partial x^i} = p_j \frac{\partial \dot{x}^j}{\partial x^i} - \frac{\partial L}{\partial x^i} - \frac{\partial L}{\partial \dot{x}^j} \frac{\partial \dot{x}^j}{\partial x^i} = -\frac{\partial L}{\partial x^i} = -\frac{d}{dt} p_i$$

and

$$\frac{\partial H}{\partial p_i} = \dot{x}^i + p_j \frac{\partial \dot{x}^j}{\partial p_i} - \frac{\partial L}{\partial \dot{x}^j} \frac{\partial \dot{x}^j}{\partial p_i} = \dot{x}^i. \quad (3)$$

So, the second order ODEs (1) in an n -dimensional space are equivalent with the first order ODEs (2) in a $2n$ -dimensional space.

The Hamiltonian H associated to L was built in 1834 by Hamilton. With this function, the Euler-Lagrange equations (1) achieve a symmetrical structure in the form of the Hamilton equations (2).

More than that, in the case when the Lagrangian L is an autonomous function (does not explicitly depend on t), the function H is a first integral for the Hamilton equations, because

$$\frac{dH}{dt} = \frac{d}{dt} (p_i \dot{x}^i - L) = \frac{dp_i}{dt} \dot{x}^i + p_i \frac{d\dot{x}^i}{dt} - \frac{\partial L}{\partial t} - \frac{\partial L}{\partial x^i} \frac{dx^i}{dt} - \frac{\partial L}{\partial \dot{x}^i} \frac{d\dot{x}^i}{dt} = -\frac{\partial L}{\partial t} = 0.$$

Using the symplectic structure $J = \begin{pmatrix} 0 & \delta_j^i \\ -\delta_j^i & 0 \end{pmatrix}$, $(\delta_j^i) \in M_n(R)$, we may write the Hamilton equations in the matrix form

$$J \begin{pmatrix} \dot{x}^j \\ \dot{p}_i \end{pmatrix} + \begin{pmatrix} \frac{\partial H}{\partial x^i} \\ \frac{\partial H}{\partial p_i} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

1.3 Multi-time Euler-Lagrange equations

We consider the multi-time variable $t = (t^1, \dots, t^p) \in R^p$, the functions $x^i : R^p \rightarrow R$, $(t^1, \dots, t^p) \rightarrow x^i(t^1, \dots, t^p)$, $i = 1, \dots, n$, and we denote $x_\alpha^i = \frac{\partial x^i}{\partial t^\alpha}$, $\alpha = 1, \dots, p$. The Lagrange function

$$L : R^{p+n+np} \rightarrow R, \quad (t^\alpha, x^i, x_\alpha^i) \rightarrow L(t^\alpha, x^i, x_\alpha^i)$$

gives the Euler-Lagrange equations

$$\frac{\partial}{\partial t^\alpha} \frac{\partial L}{\partial x_\alpha^i} = \frac{\partial L}{\partial x^i}, \quad i = 1, \dots, n, \quad \alpha = 1, \dots, p$$

(second order PDEs system on the n -dimensional space).

1.4 Multi-time Hamilton equations

In 1935 de Donder [1] obtained the Hamilton equations in the multi-time case by using the partial derivatives

$$p_k^\alpha = \frac{\partial L}{\partial x_\alpha^k} \tag{4}$$

and the Hamiltonian $H = p_k^\alpha x_\alpha^k - L$. If L satisfies some conditions, then the system (4) defines a C^1 bijective transformation $x_\alpha^i \rightarrow p_i^\alpha$, called the Legendre transformation for the multi-time case. By this transformation we have:

$$\frac{\partial H}{\partial p_i^\alpha} = x_\alpha^i + p_\beta^k \frac{\partial x_\beta^k}{\partial p_i^\alpha} - \frac{\partial L}{\partial x_\beta^k} \frac{\partial x_\beta^k}{\partial p_i^\alpha} = x_\alpha^i$$

$$\frac{\partial H}{\partial x^i} = p_k^\alpha \frac{\partial x_\alpha^k}{\partial x^i} - \frac{\partial L}{\partial x^i} - \frac{\partial L}{\partial x_\alpha^k} \frac{\partial x_\alpha^k}{\partial x^i} = -\frac{\partial L}{\partial x^i}.$$

The $np + n$ Hamilton equations $\frac{\partial x^i}{\partial t^\alpha} = \frac{\partial H}{\partial p_i^\alpha}$, $\frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{\partial H}{\partial x^i}$ (summation after α), $i = 1, \dots, n$, $\alpha = 1, \dots, p$ are first order PDEs on the space R^{n+pn} , equivalent to the Euler-Lagrange equations on R^n .

1.5 The conservation of energy-moment tensor

The multi-time Hamiltonian is not conserved on the solutions of the multi-time Hamilton equations even if the Lagrangian is autonomous (does not depend on t^α). But, we may observe that the Lagrangian L defines the energy-moment tensor

$$T_\beta^\alpha = x_\beta^i \frac{\partial L}{\partial x_\alpha^i} - \delta_\beta^\alpha L,$$

whose divergence is

$$\begin{aligned} \frac{\partial}{\partial t^\alpha} T_\beta^\alpha &= \frac{\partial}{\partial t^\alpha} (x_\beta^i p_i^\alpha) - \delta_\beta^\alpha \left(\frac{\partial L}{\partial t^\alpha} + \frac{\partial L}{\partial x^i} x_\alpha^i + \frac{\partial L}{\partial x_\gamma^i} \frac{\partial x_\gamma^i}{\partial t^\alpha} \right) = \\ &= \frac{\partial x_\beta^i}{\partial t^\alpha} p_i^\alpha + x_\beta^i \frac{\partial p_i^\alpha}{\partial t^\alpha} - \frac{\partial L}{\partial t^\beta} - \frac{\partial L}{\partial x^i} x_\beta^i - \frac{\partial L}{\partial x_\gamma^i} \frac{\partial x_\gamma^i}{\partial t^\beta} = \\ &= x_\beta^i \left(\frac{\partial p_i^\alpha}{\partial t^\alpha} - \frac{\partial L}{\partial x^i} \right) - \frac{\partial L}{\partial t^\beta} = -\frac{\partial L}{\partial t^\beta}. \end{aligned}$$

So, the energy-moment tensor T_β^α is conserved on the solutions of the multi-time Hamilton equations, if the Lagrangian L is autonomous.

2 Legendre transform of a multi-time Hamiltonian. Dual action

We may write the multi-time Hamilton equations in the form ([9],[12])

$$\delta_\beta^\alpha \delta_j^i \frac{\partial p_i^\beta}{\partial t^\alpha} + \frac{\partial H}{\partial x^j} = 0, \quad -\delta_\beta^\alpha \delta_j^i \frac{\partial x^j}{\partial t^\alpha} + \frac{\partial H}{\partial p_i^\beta} = 0$$

or

$$(\delta \otimes J) \begin{pmatrix} \frac{\partial x^j}{\partial t^\alpha} \\ \frac{\partial p_i^\beta}{\partial t^\alpha} \end{pmatrix} + \begin{pmatrix} \frac{\partial H}{\partial x^j} \\ \frac{\partial H}{\partial p_j^\beta} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (5)$$

$j = 1, \dots, n; i = 1, \dots, n; \alpha = 1, \dots, p; \beta = 1, \dots, p$, where

$$\delta \otimes J = \begin{pmatrix} 0 & \delta_\beta^\alpha \delta_j^i \\ -\delta_\beta^\alpha \delta_j^i & 0 \end{pmatrix}$$

is a polysymplectic structure acting on R^{np+np^2} with values in R^{n+np} .

Let $T_0 = [0, T^1] \times [0, T^2] \times \dots \times [0, T^p]$ and $dt = dt^1 \dots dt^p$ the volume element. The action Ψ , whose Euler-Lagrange equations are the Hamilton equations, is

$$\begin{aligned} \Psi(u) &= \int_{T_0} \mathcal{L} \left(t, u(t), \frac{\partial u}{\partial t} \right) dt, \quad \text{where } u = (x, p), \\ \mathcal{L} \left(t, u(t), \frac{\partial u}{\partial t} \right) &= -\frac{1}{2} \left(\frac{\partial p_i^\alpha}{\partial t^\alpha} x^i - \frac{\partial x^i}{\partial t^\alpha} p_i^\alpha \right) - H(t, x, p) = \\ &= -\frac{1}{2} \left(\frac{\partial p_j^\alpha}{\partial t^\alpha}, -\frac{\partial x^j}{\partial t^\beta} \right) \begin{pmatrix} \delta_{ij} & o \\ o & \delta^{\alpha\beta} \delta_{ij} \end{pmatrix} \begin{pmatrix} x^i \\ p_i^\alpha \end{pmatrix} - H(t, x(t), p(t)) = \\ &= -\frac{1}{2} G \left((\delta \otimes J) \frac{\partial u}{\partial t}, u \right) - H(t, u), \end{aligned}$$

where the scalar product is represented by the matrix $G = \begin{pmatrix} \delta_{ij} & 0 \\ 0 & \delta^{\beta\alpha} \delta_{ij} \end{pmatrix}$ (standard Riemannian metric from R^{n+np}). Indeed, the Euler-Lagrange equations produced by $\mathcal{L}$ are

$$\frac{1}{2} \frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{1}{2} \frac{\partial p_i^\alpha}{\partial t^\alpha} - \frac{\partial H}{\partial x^i}, \text{ i.e. }, \frac{\partial p_i^\alpha}{\partial t^\alpha} = -\frac{\partial H}{\partial x^i}$$

and

$$\frac{\partial x^i}{\partial t^\alpha} = \frac{\partial H}{\partial p_i^\alpha}.$$

It is well-known that the Legendre transform of a Lagrangian L is the Hamiltonian H . Besides this classical duality, we introduce a duality based on the Legendre transform of a Hamiltonian H .

The Legendre transformation $H^*(t, \cdot)$ of the Hamiltonian $H(t, \cdot)$ is defined by the implicit formula

$$H^*(t, v) = G(u, v) - H(t, u), \quad v = \nabla_u H(t, u),$$

when $\nabla H(t, u)$ is invertible. Automatically,

$$u = \nabla_v H^*(t, v),$$

so that $(\nabla H)^{-1} = \nabla H^*$. If $H(t, \cdot)$ is of class C^1 , strictly convex and has the property $\frac{H(t, u)}{|u|} \rightarrow \infty$ for $|u| \rightarrow \infty$, then $H^*(t, \cdot) \in C^1(R^{n+np}, R)$, according [2]. The norm $|u|$ is coming from the scalar product (Riemannian metric) G .

The Legendre transform $H^*(t, \cdot)$ of $H(t, \cdot)$ determines a new duality. Indeed, if we write $u = (x^i, p_i^\alpha)$, the Hamiltonian equations can be written in the compact form (5). Setting $v = (y^i, q_i^\alpha)$ defined by $y^i = -x^i$, $q_i^\alpha = -p_i^\alpha$, we obtain

$$(\delta \otimes J) \frac{\partial v}{\partial t} = \begin{pmatrix} \frac{\partial q_j^\alpha}{\partial t^\alpha} \\ -\frac{\partial y^j}{\partial t^\beta} \end{pmatrix} = \begin{pmatrix} -\frac{\partial p_j^\alpha}{\partial t^\alpha} \\ \frac{\partial x^j}{\partial t^\beta} \end{pmatrix} = -(\delta \otimes J) \frac{\partial u}{\partial t} = \nabla H(t, u).$$

or equivalently

$$u = \nabla H^*(t, (\delta \otimes J) \frac{\partial v}{\partial t}).$$

if the Legendre transform $H^*(t, \cdot)$ of $H(t, \cdot)$ exists. On the other hand

$$\begin{aligned} G\left((\delta \otimes J) \frac{\partial u}{\partial t}, u\right) &= \left(\frac{\partial p_j^\alpha}{\partial t^\alpha}, -\frac{\partial x^j}{\partial t^\beta}\right) \begin{pmatrix} \delta_{ij} & o \\ o & \delta^{\alpha\beta} \delta_{ij} \end{pmatrix} \begin{pmatrix} x^i \\ p_i^\alpha \end{pmatrix} \\ &= \frac{\partial p_i^\alpha}{\partial t^\alpha} x^i - \frac{\partial x^i}{\partial t^\alpha} p_i^\alpha \end{aligned}$$

and consequently,

$$G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v\right) = \frac{\partial q_i^\alpha}{\partial t^\alpha} y^i - \frac{\partial y^i}{\partial t^\alpha} q_i^\alpha = G\left((\delta \otimes J) \frac{\partial u}{\partial t}, u\right).$$

Using the identity

$$\begin{aligned}
\frac{1}{2}G\left(-(\delta \otimes J) \frac{\partial u}{\partial t}, u\right) &= \frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v\right) - G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v\right) \\
&= \frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v\right) - G\left((\delta \otimes J) \frac{\partial u}{\partial t}, u\right) \\
&= \frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v\right) - G\left((\delta \otimes J) \frac{\partial v}{\partial t}, u\right),
\end{aligned}$$

the action

$$\Psi(u) = \int_{T_0} [\frac{1}{2}G\left(-(\delta \otimes J) \frac{\partial u}{\partial t}, u(t)\right) - H(t, u(t))]dt.$$

can be written as

$$\Psi(u) = \int_{T_0} [\frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v(t)\right) + G\left((\delta \otimes J) \frac{\partial v}{\partial t}, u(t)\right) - H(t, u(t))]dt.$$

In this way it appears the dual action

$$\Phi(v) = \int_{T_0} [\frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v(t)\right) + H^*\left(t, (\delta \otimes J) \frac{\partial v}{\partial t}\right)]dt. \quad (6)$$

3 Extremals of dual action and multi-time Hamilton equations

In this section we describe some connections between the periodical extremals of the multi-time Hamilton dual action and the periodical solutions of the multi-time Hamilton equations.

The dual action, $\Phi(v)$, is defined on a suitable space of T-periodic functions, where $T = (T^1, \dots, T^p)$. It uses the Lagrangian

$$\mathcal{L}^*\left(t, v(t), \frac{\partial v}{\partial t}\right) = \frac{1}{2}G\left((\delta \otimes J) \frac{\partial v}{\partial t}, v(t)\right) + H^*\left(t, (\delta \otimes J) \frac{\partial v}{\partial t}\right).$$

Another useful property of the dual action is $\Phi(v + c) = \Phi(v)$, $\forall c \in R^{n+np}$. Indeed, for

$$v^* = (\delta \otimes J) \frac{\partial v}{\partial t}, \quad (7)$$

we have

$$\begin{aligned}
\Phi(v+c) &= \int_{T_0} \left[\left((\delta \otimes J) \frac{\partial(v+c)}{\partial t}, v+c \right) + H^*(t, (v+c)^*) \right] dt = \\
&= \int_{T_0} \left[\left((\delta \otimes J) \frac{\partial v}{\partial t}, v \right) + \left((\delta \otimes J) \frac{\partial v}{\partial t}, c \right) + H^*(t, v^*) \right] dt = \\
&= \int_{T_0} \left[\left((\delta \otimes J) \frac{\partial v}{\partial t}, v \right) + H^*(t, v^*) \right] dt = \Phi(v).
\end{aligned}$$

Then we may restrict the search of the critical points of $\Phi(v)$ to the set

$$\widetilde{W}_T^{1,2} = \left\{ v \in W_T^{1,2} \mid \int_{T_0} v(t) dt = 0 \right\}.$$

There are situations when it is much easier to find the critical points of the dual action Φ than those of the action Ψ . This possibility was noticed by Clark in case of single-time theory.

Between the periodical critical points of the dual action Φ and the periodical solutions of Hamilton multi-time equations, there is a close connection, as we will show in the following result.

Theorem. *Let $H : T_0 \times R^{n+np} \rightarrow R$, $(t, u) \rightarrow H(t, u)$ be a Hamiltonian, measurable in t for any $u \in R^{n+np}$ and strictly convex and continuously differentiable in u for any $t \in T_0$. Suppose there are the constants $\alpha > 0$, $\delta > 0$, and the functions $\beta, \gamma \in L^2(T_0, R^+)$ so that for any $u \in R^{n+np}$ and any $t \in T_0$, to have*

$$\frac{\delta}{2} |u|^2 - \beta(t) \leq H(t, u) \leq \frac{\alpha}{2} |u|^2 + \gamma(t). \quad (8)$$

Then, the dual action Φ is continuously differentiable on $\widetilde{W}_T^{1,2}$.

Proof. From the hypothesis, the Hamiltonian $H(t, u)$ is strictly convex in u and has the property $\frac{H(t, u)}{|u|} \rightarrow \infty$ when $|u| \rightarrow \infty$. Then it follows (see [2]) that $H^*(t, \cdot) \in C^1(R^{n+np}, R)$. From the relation (8) and the equality

$$\max_{u \in R^{n+np}} \left((v^*, u) - \frac{\alpha}{2} |u|^2 - \gamma \right) = \frac{\alpha^{-1}}{2} |v^*|^2 - \gamma, \quad (9)$$

when $\alpha > 0$, we obtain the inequalities

$$\frac{\alpha^{-1}}{2} |v^*|^2 - \gamma(t) \leq H^*(t, v^*) \leq \frac{\delta^{-1}}{2} |v^*|^2 + \beta(t) \quad (10)$$

Taking into account the Proposition 2.2 from [2, page 34], with $N = n + np$ (and $n, p = q = 2$), we obtain

$$|\nabla H^*(t, v^*)| \leq (|v^*| + \beta(t) + \gamma(t)) \delta^{-1} + 1 \leq C_1 |v^*| + C_2 (\beta(t) + \gamma(t) + 1), \quad (11)$$

where C_1, C_2 are positive constants. As $\beta + \gamma + 1 \in L^2(T_0, R^+)$, from (10) and (11), the Lagrangian

$$\mathcal{L}^* \left(t, v(t), \frac{\partial v}{\partial t} \right) = \frac{1}{2} \left((\delta \otimes J) \frac{\partial v}{\partial t}, v \right) + H^*(t, v^*)$$

satisfies conditions similar to those of Theorem 1.4 from [2, page 10]. By consequence, the dual action Φ is continuously differentiable on $W_T^{1,2}$, and on $\widetilde{W}_T^{1,2}$.

Theorem. *The same hypothesis as in the previous theorem. If $v \in \widetilde{W}_T^{1,2}$ is a critical point for the action Φ , periodical, with the period $T = (T^1, \dots, T^p)$, then the function $w(t) = \nabla H^*(t, v^*(t))$ verifies the problem*

$$(\delta \otimes J) \frac{\partial w}{\partial t} + \nabla H^*(t, w(t)) = 0, \quad w(0) = w(T)$$

and

$$v(t) = w(t) - \frac{1}{T^1 \dots T^p} \int_{T_0} w(t) dt.$$

Proof. We take $w = (z^i, r_i^\alpha)$, $v^* = (v^{*i}, v_i^{*\alpha})$. From the definition of the dual action, we have $v = (-x^i, -p_i^\alpha)$, and $v^* = \left(-\frac{\partial p_i^\alpha}{\partial t^\alpha}, \frac{\partial x^j}{\partial t^\beta} \right)$. Then

$$\mathcal{L}^* \left(t, v(t), \frac{\partial v}{\partial t} \right) = \frac{1}{2} \left(-\frac{\partial x^i}{\partial t^\alpha} p_i^\alpha + \frac{\partial p_i^\alpha}{\partial t^\alpha} x^i \right) + H^* \left(t, -\frac{\partial p_i^\alpha}{\partial t^\alpha}, \frac{\partial x^i}{\partial t^\alpha} \right).$$

If the function v is a critical point for the action produced by $\mathcal{L}^*$, then the Euler-Lagrange equations $\frac{\partial}{\partial t^\alpha} \frac{\partial \mathcal{L}^*}{\partial x_\alpha^i} = \frac{\partial \mathcal{L}^*}{\partial x^i}$, where $x_\alpha^i = \frac{\partial x^i}{\partial t^\alpha}$, are verified. So, we obtain

$$-\frac{1}{2} \frac{\partial p_i^\alpha}{\partial t^\alpha} + \frac{\partial}{\partial t^\alpha} \frac{\partial H^*}{\partial v_i^{*\alpha}} = \frac{1}{2} \frac{\partial p_i^\alpha}{\partial t^\alpha},$$

or

$$\frac{\partial}{\partial t^\alpha} \frac{\partial H^*}{\partial v_i^{*\alpha}} = \frac{\partial p_i^\alpha}{\partial t^\alpha}.$$

Because ${}^t(z^i, r_i^\alpha) = \nabla H^*(t, v^{*i}, v_i^{*\alpha})$, we find $r_i^\alpha = \frac{\partial H^*}{\partial v_i^{*\alpha}}$ and then

$$\frac{\partial r_i^\alpha}{\partial t^\alpha} = \frac{\partial p_i^\alpha}{\partial t^\alpha}. \quad (12)$$

On the other side, the equality $w = \nabla H^*\left(t, -\frac{\partial p_i^\alpha}{\partial t^\alpha}, \frac{\partial x^i}{\partial t^\alpha}\right)$ and the duality produce

$${}^t\left(-\frac{\partial p_i^\alpha}{\partial t^\alpha}, \frac{\partial x^i}{\partial t^\alpha}\right) = \nabla H(t, w).$$

By using the relation (12) we have ${}^t\left(-\frac{\partial r_i^\alpha}{\partial t^\alpha}, \frac{\partial x^i}{\partial t^\alpha}\right) = \nabla H(t, w)$, and hence

$$-\frac{\partial r_i^\alpha}{\partial t^\alpha} = \nabla_{z^i} H(t, w). \quad (13)$$

From the definition of the function H we find

$$\frac{\partial z^i}{\partial t^\alpha} = \frac{\partial H}{\partial r_i^\alpha}(t, z^i, r_i^\alpha). \quad (14)$$

From the relations (13) and (14) we obtain

$${}^t\left(-\frac{\partial r_i^\alpha}{\partial t^\alpha}, \frac{\partial z^i}{\partial t^\alpha}\right) = \nabla H(t, w) \text{ or } (\delta \otimes J) \frac{\partial w}{\partial t} = \nabla H(t, w).$$

Consequently, w is the solution of the equation $(\delta \otimes J) \frac{\partial w}{\partial t} + \nabla H^*(t, w) = 0$. We consider v a critical point for the dual action $\Phi(v)$ having mean value zero. Then, also $v_c = v + c$ is a critical point for Φ because $\int_{T_0} \left((\delta \otimes J) \frac{\partial v}{\partial t}, c \right) dt = 0$ (see the periodicity of v) and

$$\Phi(v + c) = \int_{T_0} \left((\delta \otimes J) \frac{\partial (v + c)}{\partial t}, v + c \right) + H^*(t, (v + c)^*) dt =$$

$$\begin{aligned}
&= \int_{T_0} \left((\delta \otimes J) \frac{\partial v}{\partial t}, v \right) + \left((\delta \otimes J) \frac{\partial v}{\partial t}, c \right) + H^*(t, v^*) dt = \\
&= \int_{T_0} \left((\delta \otimes J) \frac{\partial v}{\partial t}, v \right) + H^*(t, v^*) dt = \Phi(v).
\end{aligned}$$

Because v has the mean value zero, i.e., $\int_{T_0} v(t) dt = 0$, we find

$$\int_{T_0} (v_c(t) - c) dt = 0,$$

and hence

$$c = \frac{1}{T^1 \dots T^p} \int_{T_0} u_c(t) dt.$$

If v_c is a critical point for Φ , then $u_c = -v_c$ is a critical point for Ψ , i.e., it verifies the equation $(\delta \otimes J) \frac{\partial u_c}{\partial t} + \nabla H(t, u_c) = 0$. By duality,

$$u_c = \nabla H^* \left(t, {}^t(\delta \otimes J) \frac{\partial u_c}{\partial t} \right) = \nabla H^* \left(t, -(\delta \otimes J) \frac{\partial u}{\partial t} \right) = \nabla H^*(t, v^*) = w.$$

Then $v(t) = v_c(t) - c = w(t) - \frac{1}{T^1, \dots, T^p} \int_{T_0} w(t) dt$. Because v has the period $T = (T^1, \dots, T^p)$, we have $v(0) = v(T)$, so $w(0) = w(T)$.

Acknowledgements. The authors are grateful to Prof. Kostake Teleman and Ana-Maria Teleman for their valuable comments on this paper.

References

- [1] T. de Donder, *Theorie invariante du calcul des variations*, 1935.
- [2] J. Mawhin, M. Willem: *Critical Point Theory and Hamiltonian Systems*, Springer-Verlag, 1989.
- [3] M. Neagu, C. Udriște: *From PDE Systems and Metrics to Geometric Multi-time Field Theories*, Seminarul de Mecanică, Sisteme Dinamice Diferențiale, 79, Universitatea de Vest din Timișoara, 2001.
- [4] P. J. Olver, The Equivalence Problem and Canonical Forms for Quadratic Lagrangians, *Advances in Applied Mathematics*, 9 (1998), 226-257.

[5] A.-M. Teleman, C. Udriște: *On Hamiltonian Formalisms in Mathematical Physics*, 4-th International Conference of Balkan Society of Geometers, Aristotle University of Thessaloniki, 26-30 June, 2002.

[6] C. Udriște: *Nonclassical Lagrangian Dynamics and Potential Maps*, Proceedings of the Conference in Mathematics in Honour of Professor Radu Roșca on the occasion of his Ninetieth Birthday, Katholieke University Brussel, Katholieke University Leuven, Belgium, December 11-16, 1999; <http://xxx.lanl.gov/math.DS/0007060>, (2000).

[7] C. Udriște: *Geometric Dynamics*, Kluwer Academic Publishers, Dordrecht/Boston /London, Mathematics and Its Applications, 513, 2000; Southeast Asian Bulletin of Mathematics, Springer-Verlag 24 (2000).

[8] C. Udriște: *Solutions of DEs and PDEs as Potential Maps Using First Order Lagrangians*, Centenial Vranceanu, Romanian Academy, University of Bucharest, June 30-July 4, (2000); <http://xxx.lanl.gov/math.DS/0007061>, (2000); Balkan Journal of Geometry and Its Applications 6, 1, 93-108, 2001.

[9] C. Udriște: *From integral manifolds and metrics to potential maps*, Atti dell'Accademia Peloritana dei Pericolanti, Classe 1 di Scienze Fis. Mat. e Nat., 81-82, A 01006 (2003-2004), 1-14.

[10] C. Udriște, M. Neagu: *From PDE Systems and Metrics to Generalized Field Theories*, <http://xxx.lanl.gov/abs/math.DG/0101207>.

[11] C. Udriște, M. Postolache: *Atlas of Magnetic Geometric Dynamics*, Geometry Balkan Press, Bucharest, 2001.

[12] C. Udriște, A.-M. Teleman: *Hamilton Approaches of Fields Theory*, IJMMS, 57 (2004), 3045-3056; ICM Satellite Conference in Algebra and Related Topics, University of Hong-Kong, 13-18.08.02.

University Politehnica of Bucharest
Department of Mathematics
Splaiul Independentei 313
060042 Bucharest, Romania
email:udriste@mathem.pub.ro